

Lecture 10: Bias - Variance Decomposition

Concept of BIAS and VARIANCE


- Consider an example: $z_0 \leftarrow$ true location of an object

$z \leftarrow$ noisy GPS measurements of the location

treated as a random variable

it has some mean: $\mathbb{E}[z]$

it has some variance: $\mathbb{E}[(z - \mathbb{E}[z])^2]$



- Bias describes the systematic error in the measurements z (possible offset)

$$\text{BIAS: } \mathbb{E}[z] - z_0$$


- Variance describes how much the measurements vary (amount of noise in GPS measurements)

$$\text{VARIANCE: } \mathbb{E}[(z - \mathbb{E}[z])^2] = \mathbb{E}[z^2] - \mathbb{E}[z]^2$$

- Bias : $\mu_z - z_0$

Variance : $\mathbb{E}[z^2] - \mu_z^2$

- Squared error between measurement and true value : $(z - z_0)^2$

- Expected squared error : $\mathbb{E}[(z - z_0)^2]$
(Averaged)


$$\begin{aligned}&= \mathbb{E}\left[\left((z - \mu_z) + (\mu_z - z_0)\right)^2\right] \\&= \mathbb{E}\left[(z - \mu_z)^2\right] + \mathbb{E}\left[(\mu_z - z_0)^2\right] + 2 \mathbb{E}\left[(z - \mu_z)(\mu_z - z_0)\right] \\&= \underbrace{\mathbb{E}\left[(z - \mu_z)^2\right]}_{\text{variance}} + \underbrace{(\mu_z - z_0)^2}_{\text{bias}} + 2 (\mu_z - z_0) \underbrace{\left(\mathbb{E}[z] - \mu_z\right)}_{0} \quad \begin{matrix} \text{constant} & \text{constant} \\ \curvearrowleft & \curvearrowleft \\ \downarrow & \downarrow \\ \text{constant} & \text{constant} \end{matrix}\end{aligned}$$

- In other words, the averaged squared error between z and z_0 is the sum of the squared bias and variance

- To obtain a small expected squared error, we have to consider both bias and variance

- We will now apply the bias-variance concept to a regression setting
 - z_0 will now correspond to the true relationship between inputs and outputs
 - random variable z will correspond to the model learned from training data since training data T is a random collection from $p(x, y)$, the model z learned from it is also random as it is a function of training data T

- Let the true relationship between input x and output y be described by some function $f_0(x)$ plus i.i.d. noise ϵ

True model

$$y = f_0(x) + \epsilon, \quad \text{with } \mathbb{E}[\epsilon] = 0$$

similar to z_0

$$\text{Var}[\epsilon] = \sigma^2$$

- Learned model is random variable; therefore model prediction $\hat{y}(x; T)$ is r.v.!

Learned model

$$\hat{y}(x; T) \leftarrow \text{prediction from model trained on dataset } T$$

similar to z

- Let the true relationship between input \underline{x} and output y be described by some function $f_0(\underline{x})$ plus i.i.d. noise ϵ

True model

$$y = f_0(\underline{x}) + \epsilon, \quad \text{with } \mathbb{E}[\epsilon] = 0$$

\downarrow
 'z'

$$\text{Var}[\epsilon] = \sigma^2$$

- The learned model is a r.v.; therefore model prediction $\hat{y}(\underline{x}; T)$ is r.v.!
- Define average trained model corresponding to $\bar{\underline{x}}$:

$$\bar{f}(\underline{x}) = \mathbb{E}_{\mathcal{T}} [\hat{y}(\underline{x}; T)]$$

\uparrow
average model

\uparrow
expected value
over N training points
drawn from $p(\underline{x}, y)$

we would achieve if we
could re-train the model
infinite # of times on different
training datasets, each of size N

- Recall the definition of \bar{E}_{new} (for regression with squared error)

$$E_{\text{new}} = \mathbb{E}_* [(y_* - \hat{y}(\underline{x}_*; T))^2]$$

$$\bar{E}_{\text{new}} = \mathbb{E}_T [E_{\text{new}}] = \mathbb{E}_T [\mathbb{E}_* [(y_* - \hat{y}(\underline{x}_*; T))^2]]$$

- Change the order of integration

$$\bar{E}_{\text{new}} = \mathbb{E}_* [\mathbb{E}_T [(y_* - \hat{y}(\underline{x}_*; T))^2]]$$

 replace $y^* = f_o(\underline{x}^*) + \epsilon$

$$= \mathbb{E}_* [\mathbb{E}_T [(f_o(\underline{x}_*) + \epsilon - \hat{y}(\underline{x}_*; T))^2]]$$

$$= \mathbb{E}_* [\mathbb{E}_T [(\hat{y}(\underline{x}_*; T) - f_o(\underline{x}_*) - \epsilon)^2]]$$

$$= \mathbb{E}_* [\mathbb{E}_T [(\hat{y}(\underline{x}_*; T) - \bar{f}(\underline{x}_*) + \bar{f}(\underline{x}_*) - f_o(\underline{x}_*) - \epsilon)^2]]$$


 $\bar{f}(\underline{x}) = \mathbb{E}_T [\hat{y}(\underline{x}; T)]$

$$\mathbb{E}_{\text{new}} = \mathbb{E}_* \left[\mathbb{E}_T \left[\underbrace{(\hat{y}(\underline{x}_*; T) - \bar{f}(\underline{x}_*))^2}_{A_1} + \underbrace{\bar{f}(\underline{x}_*) - f_o(\underline{x}_*)}_{A_2} + \underbrace{\epsilon^2}_{A_3} \right] \right]$$

$$= \mathbb{E}_* \left[\mathbb{E}_T \left[(A_1 + A_2 + A_3)^2 \right] \right] = \mathbb{E}_* \left[\mathbb{E}_T \left[A_1^2 + A_2^2 + A_3^2 + 2(A_1 A_2 + A_2 A_3 + A_3 A_1) \right] \right]$$

$$\begin{aligned} \mathbb{E}_* \left[\mathbb{E}_T [A_1 A_2] \right] &= \mathbb{E}_* \left[\mathbb{E}_T \left[(\hat{y}(\underline{x}_*; T) - \bar{f}(\underline{x}_*)) (\bar{f}(\underline{x}_*) - f_o(\underline{x}_*)) \right] \right] \\ &= \mathbb{E}_* \left[(\bar{f}(\underline{x}_*) - f_o(\underline{x}_*)) \left(\underbrace{\mathbb{E}_T [\hat{y}(\underline{x}_*; T)] - \bar{f}(\underline{x}_*)}_{0} \right) \right] = 0 \end{aligned}$$

$$\mathbb{E}_* \left[\mathbb{E}_T [A_2 A_3] \right] = \mathbb{E}_* \left[\mathbb{E}_T [(\bar{f}(\underline{x}_*) - f_o(\underline{x}_*)) \epsilon] \right] = \mathbb{E}_* \left[f(\underline{x}_*) - f_o(\underline{x}_*) \underbrace{\mathbb{E}[\epsilon]}_0 \right] = 0$$

$$\begin{aligned} \mathbb{E}_* \left[\mathbb{E}_T [A_3 A_1] \right] &= \mathbb{E}_* \left[\mathbb{E}_T [\epsilon (\hat{y}(\underline{x}_*; T) - \bar{f}(\underline{x}_*))] \right] \quad (\text{Noise is independent of model}) \\ &= \mathbb{E}_* \left[\mathbb{E}_T [\epsilon] \right]^0 \cdot \mathbb{E}_* \left[\mathbb{E}_T [(\hat{y}(\underline{x}_*; T) - \bar{f}(\underline{x}_*))] \right] = 0 \end{aligned}$$

$$\bullet \mathbb{E}_* [\mathbb{E}_T [A_1^2]] = \mathbb{E}_* [\mathbb{E}_T [(\underbrace{\hat{y}(\underline{x}_*; T)}_{\text{'z'}} - \underbrace{\bar{f}(\underline{x}_*)}_{\text{'}\mu_z\text{'}})^2]] \quad (\mathbb{E}[(\varepsilon - \mu_\varepsilon)^2])$$

Variance

describes how much $\hat{y}(\underline{x}; T)$ varies each time

the model is trained on a different training dataset

$$\begin{aligned} \bullet \mathbb{E}_* [\mathbb{E}_T [A_2^2]] &= \mathbb{E}_* [\mathbb{E}_T [(\bar{f}(\underline{x}_*) - f_o(\underline{x}_*))^2]] \\ &= \mathbb{E}_* [(\underbrace{\bar{f}(\underline{x}_*)}_{\text{'}\mu_z\text{'}} - \underbrace{f_o(\underline{x}_*)}_{\text{'}\mu_{z_o}\text{'}})^2] \quad ((\mu_z - \mu_{z_o})^2) \end{aligned}$$

Bias²


describes how much the average trained model
 $\bar{f}(\underline{x}_*)$ differs from the true $f_o(\underline{x}_*)$

$$\bullet \mathbb{E}_* [\mathbb{E}_T [A_3^2]] = \mathbb{E}_* [\mathbb{E}_T [\epsilon^2]] = \mathbb{E}_* [\text{Var}(\epsilon) + \mu_\epsilon^2] = \mathbb{E}_* [\sigma^2] = \underbrace{\sigma^2}_{\text{Irreducible error}}$$





BIAS - VARIANCE TRADE-OFF

$$\bar{E}_{\text{new}} = \underbrace{\mathbb{E}_* \left[(\bar{f}(\underline{x}_*) - f_o(\underline{x}_*))^2 \right]}_{\text{Bias}^2} + \underbrace{\mathbb{E}_* \left[\mathbb{E}_T \left[(\hat{y}(\underline{x}_*; T) - \bar{f}(\underline{x}_*))^2 \right] \right]}_{\text{Variance}} + \underbrace{\sigma^2}_{\text{Irreducible error}}$$

- for the bias to be small, the model has to be flexible
- For the variance to be small, the model should not be very sensitive to the data points in the training set



- We also know that \bar{E}_{new} typically decreases with increasing training data



Intuitively, as the size of training data increases, we have more info about the parameters, hence the variance of prediction reduces!

Example of a simulated problem


Data Generation

- $N = 10$ data points
- Input $x \sim \text{UniformDist}(-5, 10)$
- $y = \min(0.1x^2, 3) + \epsilon$
- $\epsilon \sim \text{NormalDist}(0, 1)$


Now fit the input-output data $\{x_i, y_i\}_{i=1}^{10}$ using

- Linear regression with L_2 -regularization
- Linear regression with a quadratic polynomial and L_2 -regularization
- Linear regression with a cubic polynomial and L_2 -regularization
- Regression Tree ,
- A random forest with 10 regression trees


--- True model
 $f_0(x)$



— Mean model
 $\bar{f}(x)$




--- Different models
 $\hat{y}(x_*; T)$



learned from

different training datasets of same size

Regression tree, max depth 5



Random forest, max depth 5

